

WRONSKIANs OF THETA FUNCTIONS AND SERIES FOR $1/\pi$

ALEX BERKOVICH*, HENG HUAT CHAN, AND MICHAEL J. SCHLOSSER**

Dedicated to the memory of Professor J.M. Borwein

ABSTRACT. In this article, we define functions analogous to Ramanujan's function $f(m)$ defined in his famous paper "Modular equations and approximations to π ". We then show how these new functions lead to simpler formulas for series for $1/\pi$ associated with the classical, cubic and quartic base.

1. INTRODUCTION

Let $q = e^{\pi i \tau}$ with $\text{Im } \tau > 0$ and let

$$P(q) = 1 - 24 \sum_{k=1}^{\infty} \frac{kq^k}{1-q^k}$$

and

$$\alpha(q) = \frac{\vartheta_2^4(q)}{\vartheta_3^4(q)}, \quad (1.1)$$

where

$$\vartheta_2(q) = \sum_{k=-\infty}^{\infty} q^{(k+1/2)^2} \quad \text{and} \quad \vartheta_3(q) = \sum_{k=-\infty}^{\infty} q^{k^2}.$$

In his paper "Modular equations and approximations to π ", S. Ramanujan gave a table [18, Table III] expressing the function

$$f(\ell) := \frac{\ell P(q^{2\ell}) - P(q^2)}{\vartheta_3^2(q) \vartheta_3^2(q^\ell)} \quad (1.2)$$

in terms of $\alpha(q)$ and $\alpha(q^\ell)$ for $\ell = 2, 3, 4, 5, 7, 11, 15, 17, 19, 23, 31$ and 35 . Examples of such relations are

$$f(3) = 1 + \sqrt{\alpha(q)\alpha(q^3)} + \sqrt{(1-\alpha(q))(1-\alpha(q^3))}, \quad (1.3)$$

2010 *Mathematics Subject Classification*. Primary 11F11; Secondary 11F20, 11F27, 11Y60.

Key words and phrases. Ramanujan–Borweins series, theta functions, Dedekind eta function, modular equations, Wronskians.

*Partly supported by the Simons Foundation, Award ID: 308929.

**Partly supported by FWF Austrian Science Fund grant F50-08 within the SFB "Algorithmic and enumerative combinatorics".

and

$$f(7) = 3 \left(1 + \sqrt{\alpha(q)\alpha(q^7)} + \sqrt{(1-\alpha(q))(1-\alpha(q^7))} \right). \quad (1.4)$$

Ramanujan did not provide any proofs of these identities and until now, we are not sure how he discovered them. Ramanujan's table for $f(\ell)$ was reproduced by J.M. Borwein and P.B. Borwein in their book "Pi and the AGM" [4, p. 159, Table 5.1]. The Borweins remarked that "*The verification that $f(\ell)$ has the given form is tedious but straightforward for small ℓ . For larger ℓ , we rely on Ramanujan.*" This remark added more mysteries to Ramanujan's table of identities for $f(\ell)$.

In the paragraph after Table III of [18], Ramanujan outlined briefly the relation of these identities and his series for $1/\pi$ [18, Section 13]. A clear explanation of Ramanujan's method of deriving series for $1/\pi$ using $f(\ell)$ is now available in the book by the Borweins [4, Chapter 5].

Let

$$\eta(\tau) = q^{1/12} \prod_{k=1}^{\infty} (1 - q^{2k})$$

be the (renormalized) Dedekind η -function. It is immediate that

$$P(q^2) = 12q \frac{d\eta(\tau)}{dq} \cdot \frac{1}{\eta(\tau)},$$

and we can express $f(\ell)$ in terms of the Wronskian of $\eta(\tau)$ and $\eta(\ell\tau)$ as follows:

$$f(\ell) = \frac{12}{\eta(\tau)\eta(\ell\tau)\vartheta_3^2(q)\vartheta_3^2(q^\ell)} \det \begin{pmatrix} \eta(\tau) & \eta(\ell\tau) \\ q \frac{d\eta(\tau)}{dq} & q \frac{d\eta(\ell\tau)}{dq} \end{pmatrix}.$$

In this article, we propose to replace $f(\ell)$ by functions that are associated with Wronskians of various theta functions. For example, associated with the classical Jacobi's theta functions, we define the function

$$D_\ell(q) = \frac{1}{\vartheta_3^3(q)\vartheta_3^3(q^\ell)} \det \begin{pmatrix} \vartheta_3(q) & \vartheta_3(q^\ell) \\ q \frac{d\vartheta_3(q)}{dq} & q \frac{d\vartheta_3(q^\ell)}{dq} \end{pmatrix}.$$

The study of $D_\ell(q)$ leads us to the following general series for $1/\pi$:

Theorem 1.1. *Let $N > 1$ be an integer and let*

$$\alpha_N = \alpha(e^{-\pi\sqrt{N}}), \quad (1.5)$$

where $\alpha(q)$ is given by (1.1). Then

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(k(1 - 2\alpha_N) - \frac{2}{\sqrt{N}} D_N(e^{-\pi/\sqrt{N}}) \right) (4\alpha_N(1 - \alpha_N))^k. \quad (1.6)$$

Comparing (1.6) with the Borweins' series (5.5.13) of [4] or Theorem 2.1 of [8], we conclude that (1.6) is perhaps the simplest form of Ramanujan's series for $1/\pi$ associated with the "classical base". Using (1.6), we can derive a series for $1/\pi$ whenever

α_N and $-D_N(e^{-\pi/\sqrt{N}})$ are known. For example, we will show using modular equations satisfied by $\alpha(q)$, $\alpha(q^{13})$ and $D_{13}(q)$ that

$$\alpha_{13} = \frac{1}{2} - 3\sqrt{-18 + 5\sqrt{13}},$$

and

$$-D_{13}(e^{-\pi/\sqrt{13}}) = \frac{(-7 + 3\sqrt{13})\sqrt{-18 + 5\sqrt{13}}}{4}.$$

The series corresponding to $N = 13$ is then given by

$$\frac{1}{(6\sqrt{13}\sqrt{-18 + 5\sqrt{13}})\pi} = \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^3}{(k!)^3} \left(k + \frac{1}{4} - \frac{7}{156}\sqrt{13}\right) (649 - 180\sqrt{13})^k. \quad (1.7)$$

The identity (1.7) was given by the Borweins [4, p. 172, Table 5.2a] but since Ramanujan did not provide an expression for $f(13)$, the Borweins probably arrived at the series without using any identity associated with $f(13)$.

The article is organized as follows: In Section 2, we use the general series found by H.H. Chan, S.H. Chan and Z.G. Liu [8, Theorem 2.1] to prove Theorem 1.1. We then state a result that is an extension of [8, Theorem 2.1] and use it to derive the following analogue of (1.6):

Theorem 1.2. *Let $N > 1$ be a positive integer and α_N be given as in (1.5). Then*

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^3}{(k!)^3} \left(\frac{1 + \alpha_N}{1 - \alpha_N}k + \hat{a}_N\right) \left(-4\frac{\alpha_N}{(1 - \alpha_N)^2}\right)^k \quad (1.8)$$

where

$$\hat{a}_N = -\frac{2}{\sqrt{N}}\sqrt{\frac{\alpha_N}{1 - \alpha_N}}\hat{D}_N(e^{-\pi/\sqrt{N}}) + \frac{1}{2(1 - \alpha_N)},$$

where

$$\hat{D}_\ell(q) = D_\ell(-q).$$

Using (1.8), we derive some explicit examples, some of which are due to the Borweins. The series which we will prove with complete details is

$$\frac{1}{\sqrt{6}\pi} = \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^3}{(k!)^3} \left(\sqrt{3}(2 - \sqrt{2})k + \frac{2}{3}\sqrt{3} - \frac{5}{12}\sqrt{6}\right) (-1)^k (17 - 12\sqrt{2})^k. \quad (1.9)$$

Series (1.9) follows from the values

$$\alpha_6 = 35 + 24\sqrt{2} - 20\sqrt{3} - 14\sqrt{6},$$

and

$$\hat{D}_6(e^{-\pi/\sqrt{6}}) = \sqrt{\frac{111}{16} + 5\sqrt{2} + \frac{33}{8}\sqrt{3} + \frac{45}{16}\sqrt{6}}.$$

We observe that the terms in the sum on the right-hand side of (1.9) have alternating signs. Although series with alternating signs in the “quartic base” are present in

Ramanujan's work [18], no series with alternating signs in the “classical base” was recorded by Ramanujan. Such series should therefore be attributed to the Borweins.

In Section 3, we study the function $D_\ell(-q^2)$ and express $D_\ell(-q^2)$ in terms of Hauptmoduls when $\ell = 3, 5, 7, 11$ and 23. In Section 4, we use these identities, modular equations satisfied by $\alpha(q)$ and $\alpha(q^\ell)$, and Theorems 1.1 and 1.2 to derive several explicit series for $1/\pi$. We also provide a table of identities associated with $D_\ell(q)$ that are analogues of Ramanujan's table for $f(\ell)$. This table of formulas allows us to derive series for $1/\pi$ associated with primes other than 3, 5, 7, 11 and 23. In particular, we give an expression for $D_{13}(q)$, for which its counterpart $f(13)$ is missing in Ramanujan's table. The discovery of an expression for $D_{13}(q)$ in terms of $\alpha(q)$ and $\alpha(q^{13})$ leads to a proof of (1.7).

In Section 5, we turn our attention to the Borweins' cubic theta functions (see [5], [6]) and define the following cubic analogue of $D_\ell(q)$:

$$C_\ell(q) = \frac{1}{a^2(q)a^2(q^\ell)} \det \begin{pmatrix} a(q) & a(q^\ell) \\ q \frac{da(q)}{dq} & q \frac{da(q^\ell)}{dq} \end{pmatrix},$$

where

$$a(q) = \sum_{m,n=-\infty}^{\infty} q^{m^2+mn+n^2}.$$

We give a simpler version of Ramanujan's cubic analogue of Theorem 1.1 and a similar expression for the cubic analogue of Theorem 1.2 due to H.H. Chan, W.C. Liaw and V. Tan [13]. We then derive representations of $C_\ell(q)$ in terms of Hauptmoduls for $\ell = 2, 5$ and 11 and give several series for $1/\pi$ using these identities and modular equations satisfied by the cubic singular modulus. We end Section 5 with a table similar to that of Ramanujan for $f(\ell)$.

Define the quartic analogue of $D_\ell(q)$ as

$$Q_\ell(q) = \frac{1}{A^5(q)A^5(q^\ell)} \det \begin{pmatrix} A^2(q) & A^2(q^\ell) \\ q \frac{dA^2(q)}{dq} & q \frac{dA^2(q^\ell)}{dq} \end{pmatrix}^2,$$

where

$$A(q^2) = \frac{\eta^8(\tau) + 32\eta^8(4\tau)}{\eta^4(2\tau)}.$$

The reason why we use $A^2(q)$ instead of $A(q)$ in our definition of $Q_\ell(q)$ is because of the identity (see [2, (3.3) and (3.6)])

$$A^2(q^2) = \frac{\eta^{16}(\tau)}{\eta^8(2\tau)} \left(1 + 32 \frac{\eta^8(4\tau)}{\eta^8(\tau)} \right)^2 = \frac{\eta^{16}(\tau)}{\eta^8(2\tau)} \left(1 + 64 \frac{\eta^{24}(2\tau)}{\eta^{24}(\tau)} \right), \quad (1.10)$$

which allows us to work with modular forms associated with $\Gamma_0(2\ell) + W_\ell$ instead of $\Gamma_0(4\ell) + W_\ell$, where $\Gamma_0(N) + W_e$ denotes the group generated by $\Gamma_0(N)$ and

$$W_e = \begin{pmatrix} a\sqrt{e} & b/\sqrt{e} \\ cN/\sqrt{e} & d\sqrt{e} \end{pmatrix},$$

where $e|N$, $\gcd(N/e, e) = 1$ and $\det(W_e) = 1$. In Section 6, we state Ramanujan's quartic analogues of Theorems 1.1 and 1.2. We then give $Q_\ell(q)$ in terms of Hauptmoduls for $\ell = 3, 5, 7, 11, 23$ and give the corresponding series for $1/\pi$. We conclude the section with identities for $Q_\ell(q)$ analogous to those given by Ramanujan for $f(\ell)$.

2. NEW REPRESENTATIONS OF THE RAMANUJAN–BORWEINS SERIES FOR $1/\pi$ FOR THE “CLASSICAL BASE”

In this section, we prove Theorem 1.1 and Theorem 1.2.

Proof of Theorem 1.1. We recall from [8, (3.5)] that

$$\vartheta_3^4(q) = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} X^k(q),$$

where

$$X(q) = 4\alpha(q)(1 - \alpha(q)).$$

Using [8, Theorem 2.1], we conclude that if

$$M_N = \frac{\vartheta_3^4(q)}{\vartheta_3^4(q^N)},$$

and

$$q \frac{dX(q)}{dq} = U(q)X(q)\vartheta_3^4(q),$$

then

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} (b_N k + a_N) \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} X_N^k,$$

where

$$a_N = \frac{U\left(e^{-2\pi/\sqrt{Ns}}\right) X\left(e^{-2\pi/\sqrt{Ns}}\right) dM_N(X(q), X(q^N))}{2N \frac{dX(q)}{dq}} \Big|_{q=e^{-2\pi/\sqrt{Ns}}},$$

$$b_N = U\left(e^{-2\pi\sqrt{N/s}}\right)$$

and

$$X_N = X\left(e^{-\pi\sqrt{N}}\right).$$

It is known that [1, p. 120, Entry 9(i)]

$$q \frac{dX(q)}{dq} = (1 - 2\alpha(q))X(q)\vartheta_3^4(q). \quad (2.1)$$

This implies that

$$b_N = 1 - 2\alpha_N.$$

To prove (1.6), it remains to verify that

$$a_N = -\frac{2}{\sqrt{N}} D_N\left(e^{-\pi/\sqrt{N}}\right). \quad (2.2)$$

This follows by first observing that

$$\frac{1}{M_N} q \frac{dM_N}{dq} = -4\vartheta_3^2(q)\vartheta_3^2(q^N)D_N(q). \quad (2.3)$$

From (2.3) and (2.1), we deduce that

$$q \frac{dM_N}{dq} = \frac{dM_N}{dX} q \frac{dX}{dq} = \vartheta_3^4(q)U(q)X(q) \frac{dM_N}{dX} = -4 \frac{\vartheta_3^6(q)}{\vartheta_3^2(q^N)} D_N(q).$$

Hence,

$$U(q)X(q) \frac{dM_N(X(q), X(q^N))}{dX(q)} \Big|_{q=e^{-2\pi/\sqrt{Ns}}} = -\frac{4\vartheta_3^2(e^{-\pi/\sqrt{N}})}{\vartheta_3^2(e^{-\pi\sqrt{N}})} D_N(e^{-\pi/\sqrt{N}}).$$

We now deduce (2.2) by using the transformation formula (see for example [1, p. 43, Entry 27(ii)])

$$\vartheta_3\left(e^{-\pi/\sqrt{N}}\right) = N^{1/4}\vartheta_3\left(e^{-\pi\sqrt{N}}\right). \quad (2.4)$$

□

We now proceed to prove Theorem 1.2. We need the following generalization of [8, Theorem 2.1].

Theorem 2.1. *Let $Z(q)$ satisfy the transformation formula*

$$Z\left(e^{-2\pi/\sqrt{rs}}\right) = rZ\left(e^{-2\pi\sqrt{r/s}}\right)C\left(e^{-2\pi\sqrt{r/s}}\right), \quad (2.5)$$

where r, s are positive integers and $C(q)$ is a certain function in q . Suppose

$$Z(q) = \sum_{k=0}^{\infty} A_k X^k(q), \quad (2.6)$$

and

$$q \frac{dX(q)}{dq} = U(q)X(q)Z(q),$$

for some functions $U(q)$ and $X(q)$. Let

$$M_N(q) = \frac{Z(q)}{Z(q^N)},$$

and suppose both $M_N(q)$ and $C(q)$ can be expressed as a function of $X(q)$. Then

$$\frac{\sqrt{s}}{2\pi\sqrt{N}} = \sum_{k=0}^{\infty} (b_N k + a_N) A_k X_N^k,$$

where

$$X_N = X\left(e^{-2\pi\sqrt{N/s}}\right),$$

$$b_N = U\left(e^{-2\pi\sqrt{N/s}}\right),$$

and

$$a_N = \frac{U\left(e^{-2\pi/\sqrt{Ns}}\right)X\left(e^{-2\pi/\sqrt{Ns}}\right)}{2N} \frac{dM_N(X(q), X(q^N))}{dX(q)} \Big|_{q=e^{-2\pi/\sqrt{Ns}}}$$

$$+ \frac{U\left(e^{-2\pi\sqrt{N/s}}\right) X\left(e^{-2\pi\sqrt{N/s}}\right) \frac{dC(q)}{dX(q)} \Big|_{q=e^{-2\pi\sqrt{N/s}}}}{2C\left(e^{-2\pi\sqrt{N/s}}\right)}.$$

The above Theorem can be proved in exactly the same way as in [8]. The only difference is that we allow $Z\left(e^{-2\pi/\sqrt{Ns}}\right)$ to be expressed in terms of $Z\left(e^{-2\pi\sqrt{N/s}}\right)$ and another function $C\left(e^{-2\pi\sqrt{N/s}}\right)$. Theorem 2.1 is a generalization of Theorem 2.1 of [8] since in the latter case, the corresponding function $C(q)$ is 1.

Let

$$\vartheta_4(q) = \sum_{k=-\infty}^{\infty} (-1)^k q^{k^2}$$

and

$$Z(q) = \vartheta_4^4(q).$$

It is known from Jacobi's Triple Product identity [1, p. 37, (22.4)] that

$$\vartheta_4(q) = \frac{\eta^2(\tau/2)}{\eta(\tau)}, \quad (2.7)$$

and [1, p. 36, Entry 22])

$$\vartheta_2(q) = 2 \frac{\eta^2(2\tau)}{\eta(\tau)}. \quad (2.8)$$

Using (2.7), (2.8) and [1, p. 43, Entry 27(iii)]

$$\eta(-1/\tau) = \sqrt{-i\tau}\eta(\tau), \quad (2.9)$$

we deduce that

$$Z\left(e^{-\pi/\sqrt{N}}\right) = \vartheta_4^4\left(e^{-\pi/\sqrt{N}}\right) = N\vartheta_2^4\left(e^{-\pi\sqrt{N}}\right). \quad (2.10)$$

Using Jacobi's identity (see [1, p. 40, Entry 25(vii)] or [4, (2.1.10)])

$$\vartheta_3^4(q) = \vartheta_2^4(q) + \vartheta_4^4(q), \quad (2.11)$$

we find that

$$\vartheta_2^4\left(e^{-\pi\sqrt{N}}\right) = Z\left(e^{-\pi\sqrt{N}}\right) \frac{\vartheta_2^4\left(e^{-\pi\sqrt{N}}\right)}{\vartheta_4^4\left(e^{-\pi\sqrt{N}}\right)}. \quad (2.12)$$

Combining (2.10), (2.12) and using (1.5), we deduce that

$$Z\left(e^{-\pi/\sqrt{N}}\right) = \vartheta_4^4\left(e^{-\pi/\sqrt{N}}\right) = N Z\left(e^{-\pi\sqrt{N}}\right) \frac{\alpha_N}{1 - \alpha_N}. \quad (2.13)$$

With the above choice of $Z(q)$, we see that from the second equality of (2.12) that

$$C(q) = \frac{\vartheta_2^4(q)}{\vartheta_4^4(q)}.$$

Applying Theorem 2.1 and using (2.13), we deduce Theorem 1.2 which is a series for $1/\pi$ in terms of $\widehat{D}_\ell(q)$. The series (1.8), in a slightly different form, was discovered by the Borweins [4, p. 182, (5.5.14)].

3. THE FUNCTIONS $D_\ell(q)$ AND $\widehat{D}_\ell(q)$

In this section, instead of working with $D_\ell(q)$, we derive identities for

$$\widehat{D}_\ell(q) := D_\ell(-q) = \frac{1}{\vartheta_4^3(q)\vartheta_4^3(q^\ell)} \det \begin{pmatrix} \vartheta_4(q) & \vartheta_4(q^\ell) \\ q \frac{d\vartheta_4(q)}{dq} & q \frac{d\vartheta_4(q^\ell)}{dq} \end{pmatrix}.$$

We first establish the following fact:

Theorem 3.1. *Let ℓ be an odd prime. Then there exists some positive integer ω_ℓ depending on ℓ such that $\widehat{D}_\ell^{\omega_\ell}(q^2)$ is a modular function on $\Gamma_0(2\ell) + W_\ell$.*

Proof. Let

$$T(\tau) := \vartheta_4(q^2) = \frac{\eta^2(\tau)}{\eta(2\tau)}, \quad (3.1)$$

where the product representation of $\vartheta_4(q^2)$ follows from (2.7).

Let

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| a, b, c, d \in \mathbf{Z}, ad - bc = 1, c \equiv 0 \pmod{N} \right\}.$$

For

$$U = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(2),$$

let

$$U \circ \tau := \frac{a\tau + b}{c\tau + d}.$$

It is known, using the transformation formula of the η -function (see for example [17, p. 163] or [9, Theorem 1.2]) and (3.1), that

$$T(U \circ \tau) = \epsilon(a, b, c, d)(c\tau + d)^{1/2}T(\tau)$$

for some 24-th root of unity depending on a, b, c and d . This implies that if

$$\Psi(\tau) = \frac{1}{T(\tau)} \frac{dT}{d\tau}(\tau),$$

then

$$\Psi(U \circ \tau) = \left(\frac{c}{2(c\tau + d)} + \Psi(\tau) \right) (c\tau + d)^2. \quad (3.2)$$

Next, let ℓ be an odd prime and observe that for

$$V = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Gamma_0(2\ell),$$

$$\Psi(\ell(V \circ \tau)) = \left(\frac{\gamma}{2(\gamma\tau + \delta)} + \Psi(\ell\tau) \right) (\gamma\tau + \delta)^2. \quad (3.3)$$

Note that since

$$V \in \Gamma_0(2\ell) \subset \Gamma_0(2),$$

(3.2) also holds for the matrix V and we find that

$$S_\ell(\tau) = \Psi(\ell\tau) - \Psi(\tau)$$

is a modular form of weight 2 on $\Gamma_0(2\ell)$. It can be checked using transformation formulas for the Dedekind η -function that for some positive integer δ_ℓ ,

$$(T^2(\ell\tau)T^2(\tau))^{\delta_\ell}$$

is a modular form of weight $2\delta_\ell$ on $\Gamma_0(2\ell)$. Therefore,

$$\widehat{D}_\ell^{\delta_\ell}(q^2) = \left(\frac{S_\ell(\tau)}{T^2(\ell\tau)T^2(\tau)} \right)^{\delta_\ell}$$

is a modular function on $\Gamma_0(2\ell)$.

Next,

$$\frac{S_\ell(W_\ell \circ \tau)}{T^2(\ell(W_\ell \circ \tau))T^2(W_\ell \circ \tau)} = \pm \frac{S_\ell(\tau)}{T^2(\ell\tau)T^2(\tau)}.$$

Hence, $(\widehat{D}_\ell(q^2))^{\delta_\ell}$ is a modular function on $\Gamma_0(2\ell) + W_\ell$. In other words, ω_ℓ can be chosen to be δ_ℓ or $2\delta_\ell$. Replacing τ by $q = e^{\pi i \tau}$ and using the simple relation

$$\frac{dq}{d\tau} = \pi i q,$$

we deduce Theorem 3.1. □

We now use Theorem 3.1 to derive identities for $\widehat{D}_\ell(q)$. We first determine prime numbers ℓ for which all modular functions associated with $\Gamma_0(2\ell) + W_\ell$ are rational functions of a single function, which we shall call Hauptmodul. From the table in [12, p. 14], we find that this occurs when $\ell = 3, 5, 7, 11$ and 23. For such a prime ℓ , we construct a Hauptmodul, which is not unique, for the corresponding field of functions for $\Gamma_0(2\ell) + W_\ell$ and obtain the following identities:

Theorem 3.2. *Let*

$$H_\ell = H_\ell(\tau) = \left(\frac{\eta(2\tau)\eta(2\ell\tau)}{\eta(\tau)\eta(\ell\tau)} \right)^{\frac{24}{\ell+1}}.$$

Then

$$\widehat{D}_3(q^2) = 2H_3, \tag{3.4}$$

$$\widehat{D}_5^2(q^2) = 4H_5^2(1 + 4H_5), \tag{3.5}$$

$$\widehat{D}_7(q^2) = 2H_7(1 + 3H_7), \tag{3.6}$$

$$\widehat{D}_{11}(q^2) = 2H_{11}(1 + 4H_{11} + 5H_{11}^2), \tag{3.7}$$

$$\widehat{D}_{23}(q^2) = 2H_{23}(1 + 5H_{23} + 13H_{23}^2 + 20H_{23}^3 + 20H_{23}^4 + 11H_{23}^5). \tag{3.8}$$

Remark 3.1. Theorem 3.1 states that a certain power of $\widehat{D}_\ell(q^2)$ is in $\Gamma_0(2\ell) + W_\ell$. It is sometimes possible to express a lower power of $\widehat{D}_\ell(q^2)$ in terms of functions in $\Gamma_0(2\ell) + W_\ell$. An example of such a case is when $\ell = 7$ where we expect $\widehat{D}_7^3(q^2)$ to be a function of $\Gamma_0(14) + W_7$ instead of $\widehat{D}_7(q^2)$.

Remark 3.2. We note that since $q = e^{\pi i \tau}$, the identities given in Theorem 3.2 can all be expressed in terms of q^2 . Replacing q^2 by q , we obtain identities for $\widehat{D}_\ell(q)$ in terms of

$$H_\ell(\tau/2),$$

for $\ell = 3, 5, 7, 11$ and 23 , and these functions are in terms of infinite products with variable q . Replacing q by $-q$ and using

$$\prod_{k=1}^{\infty} (1 - (-q)^k) = \prod_{k=1}^{\infty} \frac{(1 - q^{2k})^3}{(1 - q^k)(1 - q^{4k})}, \quad (3.9)$$

we obtain identities from Theorem 3.2 expressing $D_\ell(q)$ in terms of Dedekind η -functions $\eta(\tau/2), \eta(\tau), \eta(\ell\tau/2)$ and $\eta(\ell\tau)$.

4. EXPLICIT EXAMPLES OF THEOREMS 1.1 AND 1.2

In this section, we first derive explicit series for $1/\pi$ for Theorem 3.2 for $N = 3, 5, 7, 11$ and 23 . We give complete details only for the case $N = 3$. We then derive explicit series for Theorem 1.2 for $N = 6, 10, 14, 22$ and 46 . We need to work harder in this case as our identities in Theorem 3.2 are only for $\ell = 3, 5, 7, 11$ and 23 instead of $6, 10, 14, 22$ and 46 . Again we will give complete details only for the case $N = 6$.

4.1. Case $N = 3$. Following Remark 3.2, we deduce from (3.4) that

$$D_3(q) = -2 \frac{\eta^3(\tau) \eta^3(3\tau)}{\vartheta_3^3(q) \vartheta_3^3(q^3)}, \quad (4.1)$$

where we have used (3.9) and the product representation of $\vartheta_3(q)$ [1, p. 36, Entry 22]):

$$\vartheta_3(q) = \frac{\eta^5(\tau)}{\eta^2(2\tau) \eta^2(\tau/2)}. \quad (4.2)$$

Let $\tau = i/\sqrt{3}$ in (4.1). Observe that

$$D_3(e^{-\pi/\sqrt{3}}) = -2 \frac{\eta^6(i\sqrt{3})}{\vartheta_3^6(e^{-\pi\sqrt{3}})}, \quad (4.3)$$

where we have used (2.4) and (2.9).

Next, using (2.7), (4.2) and (2.8), we immediately deduce Jacobi's identity [16, pp. 515-517]

$$\eta^{24}(\tau) = \frac{1}{28} \vartheta_3^{24}(q) \frac{\vartheta_2^8(q) \vartheta_4^8(q)}{\vartheta_3^8(q) \vartheta_3^8(q)}. \quad (4.4)$$

Letting $q = e^{-\pi\sqrt{n}}$ in (4.4), we deduce that

$$\frac{\eta^6(i\sqrt{N})}{\vartheta_3^6(e^{-\pi\sqrt{N}})} = \frac{1}{2^2} \sqrt{\alpha_N(1 - \alpha_N)}, \quad (4.5)$$

where we have used (2.11). It remains to compute α_3 . It is known that [1, p. 230, Entry 5(i)]

$$((1 - \alpha(q))(1 - \alpha(q^3)))^{1/4} + (\alpha(q)\alpha(q^3))^{1/4} = 1. \quad (4.6)$$

When $q = e^{-\pi/\sqrt{3}}$,

$$\alpha(e^{-3\pi/\sqrt{3}}) = \alpha(e^{-\pi\sqrt{3}}), \quad (4.7)$$

and

$$\alpha(e^{-\pi/\sqrt{r}}) = 1 - \alpha(e^{-\pi\sqrt{r}}), \quad (4.8)$$

with $r = 3$. Identity (4.8) is a consequence of (2.11) and (2.9). Substituting (4.7) and (4.8) into (4.6), we conclude that

$$\alpha_3(1 - \alpha_3) = \frac{1}{24},$$

which implies that

$$\alpha_3 = \frac{1}{2} - \frac{\sqrt{3}}{4}. \quad (4.9)$$

Let $n = 3$ in (4.5) and substitute (4.9) in the resulting equation to deduce that

$$D_3(e^{-\pi/\sqrt{3}}) = -\frac{1}{8},$$

and from (1.6), we deduce the following Ramanujan's series for $1/\pi$: The value we obtained from (4.3) is needed for one of Ramanujan's series from $1/\pi$. We now recall briefly how one can obtain Ramanujan's series for $1/\pi$. For more details, see [8].

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\frac{3}{2}k + \frac{1}{4}\right) \frac{1}{4^k}.$$

We have learnt from our derivation of the series corresponding to $N = 3$ that in order to derive a series for $1/\pi$ corresponding the $N = 3, 5, 7, 11, 23$ from Theorem 1.1 and Theorem 3.2, we only need the value of α_N . As such, for the following derivations of the series for $1/\pi$ corresponding to $N = 5, 7, 11, 23$ we will only discuss the evaluation of α_N .

4.2. Case $N = 5$.

The value of α_5 can be determined from the following modular equation of degree 5 [1, p. 280, Entry 13(i)]:

$$\begin{aligned} & (\alpha(q)\alpha(q^5))^{1/2} + ((1 - \alpha(q))(1 - \alpha(q^5)))^{1/2} \\ & + 2(16\alpha(q)\alpha(q^5)(1 - \alpha(q))(1 - \alpha(q^5)))^{1/6} = 1. \end{aligned}$$

This modular equation allowed us to conclude that

$$\alpha_5 = \frac{1}{2} - \sqrt{-2 + \sqrt{5}}.$$

Therefore, the series we obtained from (1.6) and Theorem 3.2 is

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\left(2\sqrt{-10 + 5\sqrt{5}}\right)k + \frac{\sqrt{-22 + 10\sqrt{5}}}{2} \right) (9 - 4\sqrt{5})^k.$$

4.3. Case $N = 7$. The value α_7 can be derived from the following modular equation of degree 7 [1, p. 314, Entry 19(i)]:

$$(\alpha(q)\alpha(q^7))^{1/8} + ((1 - \alpha(q))(1 - \alpha(q^7)))^{1/8} = 1. \quad (4.10)$$

This implies that

$$\begin{aligned} 4\alpha_7(1 - \alpha_7) &= \frac{1}{64}, \\ 1 - 2\alpha_7 &= \frac{3}{8}\sqrt{7}, \end{aligned}$$

and

$$\alpha_7 = \frac{1}{2} - \frac{3}{16}\sqrt{7}.$$

The series we obtained from (1.6) and Theorem 3.2 is

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\frac{21}{8}k + \frac{5}{16}\right) \left(\frac{1}{64}\right)^k.$$

4.4. Cases $N = 11$ and 23 .

Before we continue, we observe that our series obtained in this article depends entirely on the value of $\alpha_\ell(1 - \alpha_\ell)$ if $\Gamma_0(2\ell) + W_\ell$ has “genus 0”. The degree of the polynomial satisfied by $\alpha_\ell(1 - \alpha_\ell)$ increases in general with ℓ . In fact, if

$$\frac{\ell + 1}{8} = \frac{\nu}{s}$$

with $(\nu, s) = 1$, then $(\alpha_\ell(1 - \alpha_\ell))^{s/8}$ satisfies a polynomial equation of degree ν . For example, $(\alpha_7(1 - \alpha_7))^{1/8}$ satisfies a polynomial equation of degree 1 (see (4.10)). For $N = 11$ and 23 , we will have to solve cubic polynomial equations since $12/8 = 3/2$ and $24/8 = 3/1$.

We now continue with $N = 11$. The modular equation given by Ramanujan is [1, p. 363, Entry 7(i)]

$$\begin{aligned} &(\alpha(q)\alpha(q^{11}))^{1/4} + ((1 - \alpha(q))(1 - \alpha(q^{11})))^{1/4} \\ &+ 2(16\alpha(q)\alpha(q^{11})(1 - \alpha(q))(1 - \alpha(q^{11})))^{1/12} = 1. \end{aligned}$$

This implies that

$$\alpha_{11}(1 - \alpha_{11}) = -\frac{1}{12} \left(27 + 21\sqrt{33}\right)^{1/3} + \frac{2}{(27 + 21\sqrt{33})^{1/3}} + \frac{1}{16}.$$

The series we obtained from (1.6) and Theorem 3.2 is

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(k\sqrt{11} \cdot \sqrt{1 - \delta} + 2 \left(\delta^{1/6} - 2\delta^{1/3} + \frac{5}{4}\delta^{1/2}\right)\right) \delta^k,$$

where

$$\delta = -\frac{1}{3} \left(27 + 21\sqrt{33}\right)^{1/3} + \frac{8}{(27 + 21\sqrt{33})^{1/3}} + \frac{1}{4}.$$

A modular equation of degree 23 can be found in [1, p. 411, Entry 15(i)] and is given by

$$\begin{aligned} & (\alpha(q)\alpha(q^{23}))^{1/8} + ((1 - \alpha(q))(1 - \alpha(q^{23})))^{1/8} \\ & + 2^{2/3} (\alpha(q)\alpha(q^{23})(1 - \alpha(q))(1 - \alpha(q^{23})))^{1/24} = 1. \end{aligned}$$

From the above modular equation of degree 23, we deduce that

$$4\alpha_{23}(1 - \alpha_{23}) = \frac{1}{384} \frac{5\mu^2 - 1660 - 44\mu}{\mu},$$

where

$$\mu = \left(4724 + 924\sqrt{69}\right)^{1/3}.$$

The associated series we obtained from (1.6) and Theorem 3.2 is

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(k\sqrt{23} \cdot \sqrt{1 - X_{23}} + 2d_{23}\right) X_{23}^k,$$

where

$$d_{23} = \sqrt{2}X_{23}^{1/12} - 5X_{23}^{1/6} + \frac{13\sqrt{2}}{2}X_{23}^{1/4} - 10X_{23}^{1/3} + 5\sqrt{2}X_{23}^{5/12} - \frac{11}{4}X_{23}^{1/2}.$$

As mentioned in the beginning of this article, the Borweins remarked that to derive series for $1/\pi$ given in Theorem 1.1 corresponding to $N = 11$ and 23, we need to rely on Ramanujan's expressions for $f(11)$ and $f(23)$. We have shown here that this is not necessary and that these series can be constructed from the new identities (3.7) and (3.8).

Remark 4.1. We have cited [1] for modular equations of various degrees found by Ramanujan and used them to evaluate α_N . These modular equations are what we called Russell-type modular equations. They were studied systematically by R. Russell [19]. In fact, it is possible for us to construct Russell-type modular equations of any odd prime degree using the results found in [19]. For more details on how to compute such modular equations and its cubic analogues, see [10].

Remark 4.2. We observe that even with Russell-type modular equations, we still face the problem of obtaining roots of polynomials. For example, in the case of 11 and 23, we need to find roots of polynomials of degree 3. In other words, obtaining α_n using modular equations works only for relatively small composite or prime n . For certain n , especially those which are squarefree, we can compute α_n without using modular equations. This requires class field theory and explicit Shimura Reciprocity Law. For more details, see [7], [11], [14], [15] and [20].

We have seen how Theorem 3.2 can be used to derive explicit series for $1/\pi$. We now use these identities to derive examples for Theorem 1.2. In [4], the Borweins provided only examples to their series for even N . As such, we will first restrict our attention to the derivation of special cases of Theorem 1.2 when N is even.

Before we proceed, we observe that if ℓ is a prime then

$$\widehat{D}_{2\ell}(q)\vartheta_4^2(q)\vartheta_4^2(q^{2\ell}) = \widehat{D}_\ell(q)\vartheta_4^2(q)\vartheta_4^2(q^\ell) + \ell\widehat{D}_2(q^\ell)\vartheta_4^2(q^\ell)\vartheta_4^2(q^{2\ell}). \quad (4.11)$$

From the above, we know that we will need to derive a formula for $\widehat{D}_2(q)$ and this is given by

$$\widehat{D}_2^4(q) = \frac{1}{64^2} \frac{\alpha^4(q)}{(1 - \alpha(q))^3}. \quad (4.12)$$

The relation (4.12) can be proved by observing that both $\widehat{D}_2(q)$ and $\alpha(q)$ are modular functions corresponding to $\Gamma_0(4)$. Note that

$$\widehat{D}_2(q) \neq D_2(-q),$$

even though

$$\widehat{D}_\ell(-q) = D_\ell(-q),$$

for odd prime ℓ .

We are now ready to derive explicit series for $1/\pi$ arising from Theorem 1.2 for $N = 6, 10, 14, 22$ and 46 .

4.5. Case $N = 6$. From (4.11), we find that

$$\widehat{D}_6(e^{-\pi/\sqrt{6}}) = \widehat{D}_3(e^{-\pi/\sqrt{6}}) \frac{\vartheta_4^2(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^2(e^{-\pi\sqrt{6}})} + 3\widehat{D}_2(e^{-\pi\sqrt{\frac{3}{2}}}) \frac{\vartheta_4^2(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^2(e^{-\pi/\sqrt{6}})}. \quad (4.13)$$

In order to derive a series for $1/\pi$ using Theorem 1.2, we will need to derive the following identities:

$$\alpha_6 = 35 + 24\sqrt{2} - 20\sqrt{3} - 14\sqrt{6}, \quad (4.14)$$

$$\alpha_{2/3} = 35 - 24\sqrt{2} - 20\sqrt{3} + 14\sqrt{6}, \quad (4.15)$$

$$\widehat{D}_3^2(e^{-\pi/\sqrt{6}}) = \left(\frac{5}{2} + \frac{3}{2}\sqrt{3}\right)^2, \quad (4.16)$$

$$\widehat{D}_2^2(e^{-\pi\sqrt{\frac{3}{2}}}) = -\frac{41}{16}\sqrt{6} + \frac{99}{16} - \frac{35}{8}\sqrt{2} + \frac{29}{8}\sqrt{3}, \quad (4.17)$$

$$\frac{\vartheta_4^4(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^4(e^{-\pi/\sqrt{6}})} = 5 + \frac{8}{3}\sqrt{3} + 2\sqrt{6} + \frac{10}{3}\sqrt{2}, \quad (4.18)$$

$$\frac{\vartheta_4^4(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^4(e^{-\pi\sqrt{6}})} = -3 + 2\sqrt{2} + 2\sqrt{3} - \sqrt{6}. \quad (4.19)$$

Assuming that the above identities hold, we find that

$$\widehat{D}_6^2(e^{-\pi/\sqrt{6}}) = \left(\widehat{D}_3(e^{-\pi/\sqrt{6}}) \frac{\vartheta_4^2(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^2(e^{-\pi\sqrt{6}})} + 3\widehat{D}_2(e^{-\pi\sqrt{\frac{3}{2}}}) \frac{\vartheta_4^2(e^{-\pi\sqrt{\frac{3}{2}}})}{\vartheta_4^2(e^{-\pi/\sqrt{6}})} \right)^2$$

$$= \frac{111}{16} + 5\sqrt{2} + \frac{33}{8}\sqrt{3} + \frac{45}{16}\sqrt{6}.$$

Therefore,

$$\begin{aligned} a_6 &= \frac{2}{\sqrt{6}} \sqrt{\frac{\alpha_6}{1-\alpha_6}} \left(-\widehat{D}_6(e^{-\pi/\sqrt{6}}) \right) + \frac{1}{2(1-\alpha_6)} \\ &= -\left(\frac{1}{4} + \frac{1}{6}\sqrt{6} - \frac{1}{6}\sqrt{3} \right) + \frac{1}{2(1-\alpha_6)} \\ &= \frac{2}{3}\sqrt{3} - \frac{5}{12}\sqrt{6}. \end{aligned}$$

Note that we only need the value of α_6 to show that

$$\frac{1+\alpha_6}{1-\alpha_6} = \sqrt{3}(2-\sqrt{2}) \quad \text{and} \quad -4\frac{\alpha_6}{(1-\alpha_6)^2} = -17 + 12\sqrt{2}.$$

Hence, we obtain the following identity

$$\frac{1}{\sqrt{6}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\sqrt{3}(2-\sqrt{2})k + \frac{2}{3}\sqrt{3} - \frac{5}{12}\sqrt{6} \right) (-17 + 12\sqrt{2})^k.$$

Observe that using the product representation of $\vartheta_j(q)$, $j = 2, 3, 4$, we find that

$$-4\frac{\alpha(q)}{(1-\alpha(q))^2} = -64\frac{\eta^{24}(\tau)}{\eta^{24}(\tau/2)}. \quad (4.20)$$

We now recall the following modular equation of Ramanujan which is a consequence of [1, Chapter 17, Entry 12], namely,

$$U(\tau) + \frac{1}{U(\tau)} - 2 = V(\tau) + \frac{64}{V(\tau)} + 16, \quad (4.21)$$

where

$$U(\tau) = \left(\frac{\eta(\tau)\eta(6\tau)}{\eta(2\tau)\eta(3\tau)} \right)^{12} \quad \text{and} \quad V(\tau) = \left(\frac{\eta(\tau)\eta(3\tau)}{\eta(2\tau)\eta(6\tau)} \right)^6.$$

Substituting $\tau = i/\sqrt{6}$ in (4.21), we find, using the transformation formula for the η -function, that

$$V(i/\sqrt{6}) = 8.$$

This implies that

$$4\frac{\alpha_6}{(1-\alpha_6)^2} = 64 \left(\frac{\eta(i\sqrt{6})}{\eta(i\sqrt{3/2})} \right)^{24} = 17 - 12\sqrt{2},$$

and

$$4\frac{\alpha_{2/3}}{(1-\alpha_{2/3})^2} = 64 \left(\frac{\eta(i\sqrt{2/3})}{\eta(i/\sqrt{6})} \right)^{24} = 17 + 12\sqrt{2}.$$

This implies (4.14) and the identity (4.15).

The above method of deriving $\alpha_{2\ell}$ using $\alpha_{2/\ell}$ and modular equation is due to Ramanujan. For more details, see [18, Section 2] where α_{10} is derived.

Identity (4.16) follows from the identity

$$\begin{aligned}\widehat{D}_3^2(e^{-\pi/\sqrt{6}}) &= \frac{1}{16} \sqrt{16 \frac{\alpha_{1/6}\alpha_{3/2}}{(1-\alpha_{1/6})^2(1-\alpha_{3/2})^2}} = \frac{1}{16} \sqrt{16 \frac{(1-\alpha_6)(1-\alpha_{2/3})}{\alpha_6^2\alpha_{2/3}^2}} \\ &= \left(\frac{5}{2} + \frac{3}{2}\sqrt{3}\right)^2,\end{aligned}$$

where we have used the identity (see (4.8))

$$1 - \alpha_{1/r} = \alpha_r.$$

Identity (4.17) follows immediately from (4.12) and (4.15).

To prove (4.18) and (4.19), we observe that by (2.13), it suffices to compute

$$\frac{\vartheta_4^4(e^{-\pi\sqrt{2/3}})}{\vartheta_4^4(e^{-\pi/\sqrt{6}})}.$$

To finish this final task, we recall two identities, namely (see for example [4, 2.1.6])

$$\vartheta_3^2(q) + \vartheta_4^2(q) = 2\vartheta_3(q^2), \quad (4.22)$$

and [1, p. 214, (24.15)]

$$\frac{\vartheta_3^2(q)}{\vartheta_3^2(q^2)} = \frac{1 - \sqrt{\alpha(q^2)}}{\sqrt{1 - \alpha(q)}}. \quad (4.23)$$

From (4.22), we conclude that

$$\frac{\vartheta_4^2(q)}{\vartheta_4^2(q^2)} = \frac{1}{\sqrt{1 - \alpha(q^2)}} \left(2 - \frac{\vartheta_3^2(q)}{\vartheta_3^2(q^2)}\right) = \frac{1}{\sqrt{1 - \alpha(q^2)}} \left(2 - \frac{1 - \sqrt{\alpha(q^2)}}{\sqrt{1 - \alpha(q)}}\right).$$

This implies that

$$\begin{aligned}\frac{\vartheta_4^4(e^{-\pi/\sqrt{6}})}{\vartheta_4^4(e^{-\pi\sqrt{2/3}})} &= \frac{1}{1 - \alpha_{2/3}} \left(2 - \frac{1 - \sqrt{\alpha_{2/3}}}{\sqrt{1 - \alpha_{1/6}}}\right)^2 \\ &= -3 - 2\sqrt{2} + 2\sqrt{3} + \sqrt{6}.\end{aligned}$$

As indicated earlier, (4.18) and (4.19) follow from this computation.

4.6. $N = 10$.

We now discuss the other cases of N , namely, $N = 10, 14, 22$ and 46 . It is clear from our discussion of the case $N = 6$, to derive a series for $1/\pi$ from $\widehat{D}_N(q)$, we need, with help of the identities from Theorem 3.2, only the values for α_{2p} and $\alpha_{2/p}$. In the case of $N = 6$, we use modular equation (4.21) to derive α_6 and $\alpha_{2/3}$. We now discuss another method of deriving α_{2p} and $\alpha_{2/p}$ and we will illustrate this alternative method using the case $N = 10$. Let $\xi(q)$ be the right-hand side of (4.20), namely,

$$\xi(q) = -64 \frac{\eta^{24}(\tau)}{\eta^{24}(\tau/2)}.$$

Let $\xi_n = \xi(e^{-\pi\sqrt{n}})$. Then it can be shown that (see [7] and [11])

$$\frac{\xi_{10}}{\xi_{2/5}} + \frac{\xi_{2/5}}{\xi_{10}} = 103682.$$

Next, using (2.9), we deduce that for any positive real number n ,

$$\xi_{2n}\xi_{2/n} = 1. \quad (4.24)$$

Solving the above equation and using (4.24) for any positive integer n , we deduce that

$$\xi_{10}^2 = 51841 - 23184\sqrt{5},$$

which implies that

$$\xi_{10} = -161 + 72\sqrt{5}.$$

Using (4.20), we deduce that

$$\alpha_{10} = 323 + 144\sqrt{5} - 102\sqrt{10} - 228\sqrt{2}.$$

Similarly, we obtain

$$\alpha_{2/5} = 323 - 144\sqrt{5} - 102\sqrt{10} + 228\sqrt{2}.$$

Using these values and following what we have done for $N = 6$, we deduce that

$$\frac{1}{\sqrt{10}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\left(3\sqrt{10} - 6\sqrt{2}\right)k + \frac{23}{20}\sqrt{10} - \frac{5}{2}\sqrt{2} \right) \left(-161 + 72\sqrt{5}\right)^k.$$

4.7. $N = 14$.

The series for $1/\pi$ for $N = 14$ is not given by the Borweins. We now supply the missing series. We find that

$$\left(\frac{\xi_{14}}{\xi_{2/7}}\right)^{1/24} + \left(\frac{\xi_{2/7}}{\xi_{14}}\right)^{1/24} = 1 + \sqrt{2}.$$

This yields

$$\xi_{14} = -\left(\frac{1}{2}\sqrt{2} + \frac{1}{2} - \frac{1}{2}\sqrt{-1 + 2\sqrt{2}}\right)^{12}.$$

Using a formula of Ramanujan [3, Theorem 1.2], we deduce that

$$\alpha_{14} = \left(-2\sqrt{2} - 2 + \sqrt{8\sqrt{2} + 11}\right)^2 \left(\sqrt{10 + 8\sqrt{2}} - \sqrt{8\sqrt{2} + 11}\right)^2.$$

Similarly, we find that

$$\alpha_{2/7} = \left(-2\sqrt{2} - 2 + \sqrt{8\sqrt{2} + 11}\right)^2 \left(\sqrt{10 + 8\sqrt{2}} + \sqrt{8\sqrt{2} + 11}\right)^2.$$

From the values of α_{14} , we should expect the series for $1/\pi$ to be very complicated. We will list the algebraic numbers needed to generate the series:

$$X_{14} = -\frac{4\alpha_{14}}{(1 - \alpha_{14})^2} = -\left(\frac{1}{2}\sqrt{2} + \frac{1}{2} - \frac{1}{2}\sqrt{-1 + 2\sqrt{2}}\right)^{12},$$

$$\begin{aligned}
b_{14} &= \frac{1 + \alpha_{14}}{1 - \alpha_{14}}, \\
V_{14} &= \frac{\vartheta_4^4(e^{-\pi\sqrt{7/2}})}{\vartheta_4^4(e^{-\pi/\sqrt{14}})} \left(\widehat{D}_2(e^{-\pi\sqrt{7/2}}) \right)^2 = \frac{1}{56(1 - \alpha_{14})}, \\
U_{14} &= \frac{\vartheta_4^4(e^{-\pi\sqrt{7/2}})}{\vartheta_4^4(e^{-\pi/\sqrt{14}})} \left(\widehat{D}_7(e^{-\pi/\sqrt{14}}) \right)^2 = 4h_{14}^2(1 + 3h_{14})^2 \sqrt{\alpha_{2/7}},
\end{aligned}$$

where

$$h_{14} = \left(\frac{(1 - \alpha_{14})(1 - \alpha_{2/7})}{16^2 \alpha_{14}^2 \alpha_{2/7}^2} \right)^{1/8} = \sqrt{\frac{11}{4} + \frac{7}{4}\sqrt{2} + \frac{1}{4}\sqrt{217 + 154\sqrt{2}}}.$$

Then

$$\frac{1}{\sqrt{14}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(b_{14}k + \frac{2}{\sqrt{14}} \sqrt{\frac{\alpha_{14}}{1 - \alpha_{14}}} \left(-\sqrt{U_{14}} - 7\sqrt{V_{14}} \right) + \frac{1}{2(1 - \alpha_{14})} \right) X_{14}^k.$$

Note that in the case of $N = 14$, it is difficult to derive the series without knowing the explicit formula given by Theorem 1.2 and the corresponding identities given in Theorem 3.2. The complexity of the constants arising in this series is perhaps why the series is not given by the Borweins in their book.

4.8. $N = 22$.

Note that

$$\sqrt{\frac{\xi_{22}}{\xi_{2/11}}} + \sqrt{\frac{\xi_{2/11}}{\xi_{22}}} = 39202.$$

This yields

$$\xi_{22} = -\left(19601 - 13860\sqrt{2}\right),$$

and

$$\alpha_{22} = 39203 + 27720\sqrt{2} - 11820\sqrt{11} - 8358\sqrt{22}.$$

The corresponding series is

$$\frac{1}{2(-5\sqrt{2} + 7)\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(-33k + \frac{17\sqrt{2} - 33}{4} \right) \left(-19601 + 13860\sqrt{2} \right)^k.$$

4.9. $N = 46$.

The series for $1/\pi$ for $N = 46$ is not given by the Borweins. We can determine the values of ξ_{46} and $\xi_{2/23}$ from the following identity

$$\left(\frac{\xi_{46}}{\xi_{2/23}} \right)^{1/24} + \left(\frac{\xi_{2/23}}{\xi_{46}} \right)^{1/24} = 3 + \sqrt{2}.$$

We find that

$$\xi_{46} = - \left(\frac{3}{2} + \frac{\sqrt{2}}{2} - \frac{1}{2} \sqrt{7 + 6\sqrt{2}} \right)^{12}.$$

Therefore,

$$\alpha_{46} = \left(26 + 18\sqrt{2} - 3\sqrt{147 + 104\sqrt{2}} \right)^2 \left(\sqrt{1332 + 936\sqrt{2}} - 3\sqrt{147 + 104\sqrt{2}} \right)^2,$$

and

$$\alpha_{2/23} = \left(26 + 18\sqrt{2} - 3\sqrt{147 + 104\sqrt{2}} \right)^2 \left(\sqrt{1332 + 936\sqrt{2}} + 3\sqrt{147 + 104\sqrt{2}} \right)^2.$$

The following constants will give rise to an explicit series for $1/\pi$ associated with $N = 46$:

$$\begin{aligned} b_{46} &= 78\sqrt{147 + 104\sqrt{2}} + 54\sqrt{2}\sqrt{147 + 104\sqrt{2}} \\ &\quad - 3\sqrt{2}\sqrt{147 + 104\sqrt{2}}\sqrt{661 + 468\sqrt{2}}, \\ X_{46} &= - \left(\frac{3}{2} + \frac{\sqrt{2}}{2} - \frac{1}{2} \sqrt{7 + 6\sqrt{2}} \right)^{12}, \\ V_{46} &= \frac{1}{2^3 \cdot 23(1 - \alpha_{46})}, \\ U_{46} &= (2h_{46} (1 + 5h_{46} + 13h_{46}^2 + 20h_{46}^3 + 20h_{46}^4 + 11h_{46}^5))^2 \sqrt{\alpha_{2/23}}, \end{aligned}$$

where

$$h_{46} = \left(\frac{(1 - \alpha_{46})(1 - \alpha_{2/23})}{16^2 \alpha_{46}^2 \alpha_{2/23}^2} \right)^{1/24}.$$

Then

$$\frac{1}{\sqrt{46}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(b_{46}k + \frac{2}{\sqrt{46}} \sqrt{\frac{\alpha_{46}}{1 - \alpha_{46}}} \left(-\sqrt{U_{46}} - 23\sqrt{V_{46}} \right) + \frac{1}{2(1 - \alpha_{46})} \right) X_{46}^k.$$

We have, in our attempt to prove some of Borweins identities [4, p. 172, Tables 5.2a, 5.2b], used (1.6) to derive series for $1/\pi$ when N is odd and (1.8) when N is even. We would like to emphasize here that these restrictions are not necessary. Indeed if we consider $N = 6, 10$ and 22 , we obtain from (1.6) the following series for $1/\pi$:

The identity

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} (b_N k + a_N) X_N^k \quad (4.25)$$

is true when

$$\begin{aligned} b_6 &= -69 - 48\sqrt{2} + 40\sqrt{3} + 28\sqrt{6}, \\ a_6 &= -30 - 21\sqrt{2} + \frac{52}{3}\sqrt{3} + \frac{73}{6}\sqrt{6}, \end{aligned}$$

$$\begin{aligned}
X_6 &= -18872 - 13344\sqrt{2} + 10896\sqrt{3} + 7704\sqrt{6}, \\
b_{10} &= -645 + 456\sqrt{2} + 204\sqrt{10} - 288\sqrt{5}, \\
a_{10} &= -290 + 205\sqrt{2} + \frac{917}{10}\sqrt{10} - \frac{648}{5}\sqrt{5}, \\
X_{10} &= -1662776 + 1175760\sqrt{2} + 525816\sqrt{10} - 743616\sqrt{5},
\end{aligned}$$

and

$$\begin{aligned}
b_{22} &= -78405 - 55440\sqrt{2} + 23640\sqrt{11} + 16716\sqrt{22}, \\
a_{22} &= -36542 - 25839\sqrt{2} + \frac{121196}{11}\sqrt{11} + \frac{171397}{22}\sqrt{22}, \\
X_{22} &= -24589219256 - 17387203680\sqrt{2} + 7413928560\sqrt{11} + 5242439160\sqrt{22}.
\end{aligned}$$

Similarly, we also found series associated with (1.8) when N is odd. For example, when $N = 3$ and 7 , we have relatively simple series which we missed so far. These are respectively

$$\frac{1}{\sqrt{3}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\left(\sqrt{15} - 8\sqrt{3} \right) k + 6 - \frac{10}{3}\sqrt{3} \right) \left(-416 + 240\sqrt{3} \right)^k,$$

and

$$\frac{1}{\sqrt{7}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\left(255 - 96\sqrt{7} \right) k + 112 - \frac{296}{7}\sqrt{7} \right) \left(-129536 + 48960\sqrt{7} \right)^k.$$

There is also a series for the case $N = 5$ and it is given by

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\hat{b}_N k + \hat{a}_N \right) \hat{X}_N^k,$$

where

$$\begin{aligned}
\hat{b}_5 &= 35 + 16\sqrt{5} - 72\sqrt{\sqrt{5} - 2} - 32\sqrt{5\sqrt{5} - 10}, \\
\hat{a}_5 &= 15 + \frac{34}{5}\sqrt{5} - 18\sqrt{\sqrt{5} - 2} - 8\sqrt{5\sqrt{5} - 10} - \frac{1}{5}\sqrt{1990 + 890\sqrt{5}}, \\
\hat{X}_5 &= -4936 - 2208\sqrt{5} + 10160\sqrt{\sqrt{5} - 2} + 4544\sqrt{5\sqrt{5} - 10}.
\end{aligned}$$

We note that identities such as those given in Theorem 3.2 exist only when $\Gamma_0(2\ell) + W_\ell$ has genus 0 or according to [12, p. 14], when $\ell = 3, 5, 7, 11, 23$. In order to compute $D_\ell(q)$ for primes other than 3, 5, 7, 11 and 23, we introduce modular functions similar to those used by Ramanujan in his representations of $f(\ell)$.

In the following table, we will state the value of N and in each entry, set

$$\beta = \alpha(q^N).$$

Table 1: Table of identities for $D_N(q)$

	$N = 3$
	$D_3(q) = -\frac{(\alpha\beta(1-\alpha)(1-\beta))^{1/4}}{2}.$
Let	$N = 5$
	$X = \frac{(2^{10}\alpha\beta(1-\alpha)(1-\beta))^{1/6}}{8},$
then	$D_5^2(q) = 4X^2(1-4X).$
Let	$N = 7$
	$X = \frac{(\alpha\beta(1-\alpha)(1-\beta))^{1/8}}{2},$
then	$D_7(q) = -2X(1-3X).$
Let	$N = 11$
	$X = \frac{(2^4\alpha\beta(1-\alpha)(1-\beta))^{1/12}}{2},$
then	$D_{11}(q) = -2X(1-4X+5X^2).$
Let	$N = 13$
	$X = \frac{(\alpha\beta(1-\alpha)(1-\beta))^{1/2}}{16},$
and	
	$Y = \frac{1 - (\alpha\beta)^{1/2} - ((1-\alpha)(1-\beta))^{1/2}}{8},$
then	
	$10XD_{13}^4(q) + (-116X - 404XY^2 + 528XY - Y^2 + Y^3 + 1280X^2)D_{13}^2(q)$ $- 16X - 20Y^5 - 16000X^2Y - 176XY^2 + 2112X^2 + 4Y^4$ $+ 37824X^2Y^2 - 3504XY^3 + 8240XY^4 - 23040X^3 + 192XY = 0.$

$$N = 17$$

Let

$$X = \frac{(2^4 \alpha \beta (1 - \alpha)(1 - \beta))^{1/6}}{4},$$

and

$$Y = \frac{1 - (\alpha \beta)^{1/2} - ((1 - \alpha)(1 - \beta))^{1/2}}{8},$$

then

$$D_{17}^2(q) = 4 \frac{64X^3Y - 11X^2Y - 4X^2 - 24XY + 31XY^2 - 32X^3 + Y^2 + 3X - 8Y^2}{1 - Y + 5X}.$$

$$N = 19$$

Let

$$X = \frac{(\alpha \beta (1 - \alpha)(1 - \beta))^{1/4}}{4},$$

and

$$Y = \frac{1 - (\alpha \beta)^{1/4} - ((1 - \alpha)(1 - \beta))^{1/4}}{4},$$

then

$$D_{19}(q) = \frac{2X + 2Y + 16Y^3 - 10Y - 18XY}{Y - 1}.$$

$$N = 23$$

Let $X = \frac{(2^{16} \alpha \beta (1 - \alpha)(1 - \beta))^{1/24}}{2}$, then

$$D_{23}(q) = -2X(1 - 5X + 13X^2 - 20X^3 + 20X^4 - 11X^5).$$

$$N = 29$$

Let

$$X = \left(\frac{\alpha \beta (1 - \alpha)(1 - \beta)}{256} \right)^{1/6},$$

and

$$Y = \frac{1 - (\alpha \beta)^{1/2} - ((1 - \alpha)(1 - \beta))^{1/2}}{8},$$

then

$$A_2 D_{29}^4(q) + A_1 D_{29}^2(q) + A_0 = 0,$$

where

$$A_2 = -585689508612X^2,$$

$$\begin{aligned}
A_1 &= 123736544264XY + 3702335691264X^2 + 97491959398X \\
&\quad + 134904595824360X^4 - 44395652981864X^2Y^2 - 432321617914Y^3 \\
&\quad - 42626822690432X^5 - 29875947341036X^3 - 9779263696654XY^2 \\
&\quad + 41705207079730X^2Y - 8251360353152X^4Y + 5451791661904XY^3 \\
&\quad - 176409878302552X^3Y, \\
A_0 &= 13753900119256887X^2Y^3 - 4877930791543X \\
&\quad + 2618284012843192X^3Y + 2305243907550368XY^3 \\
&\quad - 700939761749206XY^2 + 505394444931798X^2Y \\
&\quad + 96399537859592XY - 4086296883979928X^2Y^2 \\
&\quad - 14225126607270367X^3Y^2 + 4709822410848252X^5Y \\
&\quad + 25397795278722548X^3Y^3 - 16793873356376932X^4Y^2 \\
&\quad + 1068896146837092XY^5 - 175149710486642X^3 \\
&\quad + 5709212469240785X^4Y - 3216747114074433XY^4 \\
&\quad - 16394572380315964X^2Y^4 - 1065063721978775X^5 \\
&\quad - 368335914073064X^4 - 19806094957276X^2 \\
&\quad + 6607217263199Y^5 - 204688220825404X^6 - 25269791081528Y^6.
\end{aligned}$$

$$N = 31$$

Let

$$X = \frac{(\alpha\beta(1-\alpha)(1-\beta))^{1/8}}{2},$$

and

$$Y = \frac{1 - (\alpha\beta)^{1/8} - ((1-\alpha)(1-\beta))^{1/8}}{8},$$

then

$$D_{31}(q) = -82X^2 + 22X - 1536Y^3 - 8Y - 32XY + 160Y^2 + 896XY^2.$$

Using the identity associated with $D_{29}(q)$, we obtain the Borweins' series [4, p. 172] associated with $N = 58$ in Theorem 1.2, namely,

$$\frac{1}{\sqrt{N}\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(k!)^3} \left(\widehat{b}_N k + \widehat{a}_N\right) \widehat{X}_N^k,$$

where

$$\widehat{b}_{58} = -6930\sqrt{2} + 1287\sqrt{58},$$

$$\begin{aligned}\widehat{a}_{58} &= -\frac{6351}{2}\sqrt{2} + \frac{68403}{116}\sqrt{58}, \\ \widehat{X}_{58} &= -192119201 + 35675640\sqrt{29}.\end{aligned}$$

As in the case of $N = 13$ for Theorem 1.1, the Borweins derived the above series without the knowledge of Ramanujan's $f(29)$, which is not listed in Ramanujan's table for $f(\ell)$.

Remark 4.3. Note that the above table contains identities analogous to Ramanujan's table for $f(\ell)$. Using the identity for $D_{13}(q)$, we obtain the series (1.7). These identities are found by computer algebra, more specifically F.G. Garvan's QSERIES package (available at <http://www.qseries.org/fgarvan/qmaple/qseries/>), using suitable functions such as X and Y given in the table. Once an identity is found, we verify the validity of the identity by deriving a modular equation from this identity and verify the modular equation using the usual technique of comparing q -series expansions of the modular functions involved in the modular equation. The identity we wish to prove is then one of the solutions of the modular equation.

5. SERIES FOR $1/\pi$ ASSOCIATED WITH THE CUBIC THETA FUNCTION $a(q)$

In this section, we consider the cubic analogue of (1.6) and (1.8). Let

$$a(q) = \sum_{m=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} q^{m^2+mn+n^2},$$

and

$$\frac{1}{\alpha^\dagger(q^2)} = 1 + \frac{1}{27} \frac{\eta^{12}(\tau)}{\eta^{12}(3\tau)}.$$

The analogue of Theorems 1.1 and 1.2 are respectively given as follows:

Theorem 5.1. *Let N be a positive integer,*

$$\alpha_N^\dagger = \alpha^\dagger\left(e^{-2\pi\sqrt{N/3}}\right),$$

and

$$C_N(q) = \frac{1}{a^2(q)a^2(q^N)} \det \begin{pmatrix} a(q) & a(q^N) \\ q \frac{da(q)}{dq} & q \frac{da(q^N)}{dq} \end{pmatrix}.$$

Then

$$\sqrt{\frac{3}{N}} \frac{1}{2\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(b_N^\dagger k + a_N^\dagger\right) \left(X_N^\dagger\right)^k, \quad (5.1)$$

where

$$\begin{aligned}b_N^\dagger &= 1 - 2\alpha_N^\dagger, \\ a_N^\dagger &= -\frac{C_N(q)}{\sqrt{N}} \Big|_{q=e^{-2\pi/\sqrt{3N}}},\end{aligned}$$

and

$$X_N^\dagger = 4\alpha_N^\dagger (1 - \alpha_N^\dagger).$$

Theorem 5.2. *Let N be a positive integer,*

$$\hat{\alpha}_N^\dagger = \alpha^\dagger \left(-e^{-\pi\sqrt{N/3}} \right),$$

and

$$\hat{C}_N(q) = \frac{1}{a^2(-q)a^2(-q^N)} \det \begin{pmatrix} a(-q) & a(-q^N) \\ q \frac{da(-q)}{dq} & q \frac{da(-q^N)}{dq} \end{pmatrix}.$$

Then

$$\sqrt{\frac{3}{N}} \frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(\hat{b}_N^\dagger k + \hat{a}_N^\dagger \right) \left(\hat{X}_N^\dagger \right)^k, \quad (5.2)$$

where

$$\begin{aligned} \hat{b}_N^\dagger &= 1 - 2\hat{\alpha}_N^\dagger, \\ \hat{a}_N^\dagger &= \frac{\hat{C}_N(q)}{\sqrt{N}} \Big|_{q=e^{-\pi/\sqrt{3N}}}, \end{aligned}$$

and

$$\hat{X}_N^\dagger = 4\hat{\alpha}_N^\dagger (1 - \hat{\alpha}_N^\dagger).$$

We now state a few identities similar to those in Theorem 3.2 for the cubic case.

Theorem 5.3. *The following holds:*

Let

$$H_\ell^\dagger = H_\ell^\dagger(\tau) = \left(\frac{\eta(3\tau)\eta(3\ell\tau)}{\eta(\tau)\eta(\ell\tau)} \right)^{\frac{12}{\ell+1}}.$$

Then

$$\begin{aligned} C_2(q^2) &= -6 \frac{H_2^\dagger}{(9H_2^\dagger + 1)^2}, \\ C_5(q^2) &= -6H_5^\dagger \frac{4H_5^\dagger + 1 + 9H_5^{\dagger 2}}{(9H_5^{\dagger 2} + 9H_5^\dagger + 1)^2}, \\ C_{11}(q^2) &= -6H_{11}^\dagger(\tau) \frac{U(H_{11}^\dagger(\tau))}{V^2(H_{11}^\dagger(\tau))}, \end{aligned} \quad (5.3)$$

where

$$U(s) = 1 + 5s + 18s^2 + 37s^3 + 54s^4 + 45s^5 + 27s^6,$$

and

$$V(s) = 1 + 9s + 18s^2 + 27s^3 + 9s^4.$$

The examples of (5.1) which follow from Theorem 5.3 are given as follows:

5.1. **Case $N = 2$.**

When $N = 2$, $\alpha_2^\dagger = \frac{\sqrt{2}-1}{2\sqrt{2}}$, $b_2^\dagger = \frac{\sqrt{2}}{2}$ and $X_2^\dagger = \frac{1}{2}$. Using (5.3) and the fact that

$$H_2^\dagger(i/\sqrt{6}) = \frac{1}{9},$$

we deduce that

$$C_2(e^{-2\pi/\sqrt{6}}) = -\frac{1}{6}.$$

The series for $1/\pi$ in this case is

$$\frac{\sqrt{3}}{2\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} (6k+1) \frac{1}{2^k}.$$

For case $N = 5$ and 11, identity (5.1) holds for the following values:

$$b_5^\dagger = \frac{11}{23}\sqrt{5}, \quad a_5^\dagger = \frac{4}{75}\sqrt{5}, \quad X_5^\dagger = \frac{4}{125},$$

and

$$\begin{aligned} b_{11}^\dagger &= -\frac{5}{242}\sqrt{11} + \frac{44}{242}\sqrt{33}, \\ a_{11}^\dagger &= -\frac{13}{726}\sqrt{11} + \frac{3}{121}\sqrt{33}, \\ X_{11}^\dagger &= -\frac{194}{1331} + \frac{225}{2662}\sqrt{3}. \end{aligned}$$

There is only one series which is a special case of (5.2) arising from Theorem 5.3 because the series on the right-hand side of (5.2) diverges for $N = 2$ and 3. This series corresponds to $N = 11$ and is given by

$$\frac{\sqrt{3}}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(\left(\frac{45}{22}\sqrt{3} + \frac{5}{22} \right) k + \frac{13}{66} + \frac{3}{11}\sqrt{3} \right) \left(-\frac{194}{1331} - \frac{225}{2662}\sqrt{3} \right)^k.$$

We end this section with cubic analogues of Ramanujan's identities for $f(\ell)$. In the following table, we will state the value of N and in each entry, set

$$\beta^\dagger = \alpha^\dagger(q^N).$$

Table 2: Table of identities for $C_N(q)$

$N = 2$

Let

$$X = \frac{(\alpha^\dagger \beta^\dagger (1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/3}}{9},$$

then

$$C_2(q) = -6X.$$

$$N = 5$$

Let

$$X = \frac{(\alpha^\dagger \beta^\dagger (1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/6}}{3},$$

then

$$C_5(q) = -6X(1 - 5X).$$

$$N = 11$$

Let

$$X = \frac{(\alpha^\dagger \beta^\dagger (1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/6}}{3},$$

and

$$Y = \frac{1 - (\alpha^\dagger \beta^\dagger)^{1/3} - ((1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/3}}{9},$$

then

$$C_{11}(q) = -33XY + 3X - 6Y + 33Y^2.$$

$$N = 17$$

Let

$$X = \frac{(\alpha^\dagger \beta^\dagger (1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/6}}{3},$$

and

$$Y = \frac{1 - (\alpha^\dagger \beta^\dagger)^{1/3} - ((1 - \alpha^\dagger)(1 - \beta^\dagger))^{1/3}}{9},$$

then

$$C_{17}(q) = \frac{6(-2X^2 + 34X^2Y + 51XY^2 - 14Y^2 - 9XY + Y + 51Y^3)}{8Y - 1}.$$

Remark 5.1. In the above table, we give an identity for $\ell = 17$ to illustrate the fact that we can compute $C_\ell(q)$ even when the genus of $\Gamma_0(3\ell) + W_\ell$ is not zero.

6. QUARTIC THEORY

In this section, we discuss series for $1/\pi$ corresponding to the quartic theory. Let

$$A^2(q^2) = \frac{\eta^{16}(\tau)}{\eta^8(2\tau)} \left(1 + 32 \frac{\eta^8(4\tau)}{\eta^8(\tau)} \right)^2 = \frac{\eta^{16}(\tau)}{\eta^8(2\tau)} \left(1 + 64 \frac{\eta^{24}(2\tau)}{\eta^{24}(\tau)} \right),$$

and

$$\frac{1}{\alpha^\perp(q^2)} = 1 + \frac{1}{64} \frac{\eta^{24}(\tau)}{\eta^{24}(2\tau)}.$$

The quartic analogues of Theorem 1.1 and 1.2 are given respectively as follows:

Theorem 6.1. *Let*

$$\alpha_N^\perp = \alpha^\perp \left(e^{-\pi\sqrt{2N}} \right),$$

and

$$Q_N(q) = \frac{1}{A^5(q)A^5(q^N)} \det \begin{pmatrix} A^2(q) & A^2(q^N) \\ q \frac{dA^2(q)}{dq} & q \frac{dA^2(q^N)}{dq} \end{pmatrix}^2.$$

Then

$$\sqrt{\frac{2}{N}} \frac{1}{2\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(b_N^\perp k + a_N^\perp \right) \left(X_N^\perp \right)^k, \quad (6.1)$$

where

$$\begin{aligned} b_N^\perp &= 1 - 2\alpha_N^\perp, \\ a_N^\perp &= \frac{\sqrt{Q_N(q)}}{4\sqrt{N}} \Big|_{q=e^{-2\pi/\sqrt{2N}}}, \end{aligned}$$

and

$$X_N^\perp = 4\alpha_N^\perp \left(1 - \alpha_N^\perp \right),$$

Theorem 6.2. *Let*

$$\begin{aligned} \hat{\alpha}^\perp(q) &= \alpha^\perp(-q), \\ \hat{\alpha}_N^\perp &= \hat{\alpha}^\perp \left(e^{-\pi\sqrt{N}} \right), \end{aligned}$$

and

$$\hat{Q}_N(q) = Q_N(-q).$$

Then

$$\sqrt{\frac{1}{N}} \frac{1}{\pi} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(\hat{b}_N^\perp k + \hat{a}_N^\perp \right) \left(\hat{X}_N^\perp \right)^k, \quad (6.2)$$

where

$$\hat{a}_N^\perp = \frac{\sqrt{|\hat{Q}_N(q)|}}{4\sqrt{N}} \Big|_{q=e^{-\pi/\sqrt{N}}},$$

$$\widehat{b}_N^\perp = 1 - 2\widehat{\alpha}_N^\perp,$$

and

$$\widehat{X}_N^\perp = 4\widehat{\alpha}_N^\perp (1 - \widehat{\alpha}_N^\perp).$$

Next, we state the quartic analogues of Theorem 3.2:

Theorem 6.3. *Let H_ℓ with $\ell = 3, 5, 7, 11$ and 13 be defined as in Theorem 3.2. Then*

$$\begin{aligned} Q_3(q^2) &= 2304 \frac{H_3^2(1 + 8H_3)^2}{(32H_3 + 1 + 64H_3^2)^3}, \\ Q_5(q^2) &= 2304 \frac{(4H_5 + 1)^2 H_5^2}{(1 + 28H_5 + 16H_5^2)^3}, \\ Q_7(q^2) &= 2304 \frac{H_7^2(1 + 11H_7 + 48H_7^2 + 88H_7^3 + 64H_7^4)^2}{(64H_7^4 + 1 + 32H_7 + 160H_7^2 + 256H_7^3)^3}, \\ Q_{11}(q^2) &= 2304 H_{11}^2 \frac{U_1^2(H_{11})}{V_1^3(H_{11})}, \\ Q_{23}(q^2) &= 2304 \frac{U_2^2(H_{23})}{V_2^3(H_{23})}, \end{aligned}$$

where

$$\begin{aligned} U_1(s) &= 1 + 12s + 69s^2 + 224s^3 + 448s^4 + 552s^5 + 384s^6 + 128s^7, \\ V_1(s) &= 1 + 32s + 192s^2 + 544s^3 + 768s^4 + 512s^5 + 64s^6, \\ U_2(s) &= 48s + 624s^2 + 4464s^3 + 21312s^4 + 75456s^5 + 209808s^6 + 472320s^7 \\ &\quad + 877056s^8 + 1357056s^9 + 1754112s^{10} + 1889280s^{11} + 1678464s^{12} \\ &\quad + 1207296s^{13} + 681984s^{14} + 285696s^{15} + 79872s^{16} + 12288s^{17}, \end{aligned}$$

and

$$\begin{aligned} V_2(s) &= 1 + 32s + 224s^2 + 928s^3 + 2624s^4 + 5504s^5 + 8864s^6 \\ &\quad + 11008s^7 + 10496s^8 + 7424s^9 + 3584s^{10} + 1024s^{11} + 64s^{12}. \end{aligned}$$

In order to compute explicit examples corresponding to (6.1), we need a way to compute $\alpha_N^\perp$. Fortunately, from the second equality of (1.10) [2, (3.6)] and the definitions of $\alpha(q)$ and $\alpha^\perp(q)$, we find that

$$\alpha^\perp(q) = \frac{4\alpha(q)}{(1 + \alpha^2(q))^2}. \quad (6.3)$$

Write $\alpha_N^\perp = \alpha^\perp(e^{-\pi\sqrt{2N}})$ and, as usual, write $\alpha_N = \alpha(e^{-\pi\sqrt{N}})$. With the above notation, we deduce from (6.3) that

$$\alpha_N^\perp = \frac{4\alpha_{2N}}{(1 + \alpha_{2N})^2}.$$

This implies that the value of $\alpha_N^\perp$ can be deduced from the value of α_{2N} , which we have already computed in Section 3 for $N = 3, 5, 7, 11$ and 23. With these values, together with Theorems 6.1 and 6.3, we obtain the following series for $1/\pi$:

Identity (6.1) is true when

$$\begin{aligned} b_3^\perp &= \frac{2\sqrt{2}}{3}, \quad a_3^\perp = \frac{\sqrt{2}}{12}, \quad X_3^\perp = \frac{1}{9}, \\ b_5^\perp &= \frac{4}{9}\sqrt{5}, \quad a_5^\perp = \frac{2}{45}\sqrt{5}, \quad X_5^\perp = \frac{1}{81}, \\ b_7^\perp &= \frac{2}{7}\sqrt{-50 + 44\sqrt{2}}, \\ a_7^\perp &= \frac{3}{196}\sqrt{-154 + 112\sqrt{2}\sqrt{7}}, \\ X_7^\perp &= \frac{249}{49} - \frac{176}{49}\sqrt{2}, \\ b_{11}^\perp &= \frac{70}{99}\sqrt{2}, \quad a_{11}^\perp = \frac{19}{396}\sqrt{2}, \quad X_{11}^\perp = \frac{1}{9801}, \end{aligned}$$

and

$$\begin{aligned} b_{23}^\perp &= \frac{14}{207}\sqrt{2}\sqrt{-1 + 78\sqrt{2}}, \\ a_{23}^\perp &= \frac{1}{828}\sqrt{2}\sqrt{-8443 + 6504\sqrt{2}}, \\ X_{23}^\perp &= \frac{43241}{42849} - \frac{10192}{14283}\sqrt{2}. \end{aligned}$$

Using values of $\hat{\alpha}_N^\perp$, Theorems 6.2 and 6.3, we deduce that (6.2) is true when

$$\hat{b}_5^\perp = \frac{\sqrt{5}}{2}, \quad \hat{a}_5^\perp = \frac{3\sqrt{5}}{40}, \quad \hat{X}_5^\perp = -\frac{1}{4},$$

and

$$\hat{b}_7^\perp = \frac{65}{63}, \quad \hat{a}_7^\perp = \frac{8}{63}, \quad \hat{X}_7^\perp = -\frac{256}{3969}.$$

For the cases $N = 11$ and $N = 23$, the values are complicated. We state them as follows: Let

$$\rho = 66599 + 11319\sqrt{33}, \quad \delta = 1389961 + 357357\sqrt{33}, \quad \text{and} \quad \epsilon = 48978 + 8526\sqrt{33},$$

then (6.2) is true when

$$\begin{aligned} \hat{b}_{11}^\perp &= -\frac{377}{1617} + \frac{32}{1617}\rho^{1/3} + \frac{665599}{17709384}\rho^{2/3} - \frac{7}{10952}\sqrt{33}\rho^{2/3}, \\ \hat{a}_{11}^\perp &= \frac{2}{539}\epsilon^{1/3} + \frac{2721}{8624}\epsilon^{2/3} - \frac{29}{176}\sqrt{33}\epsilon^{2/3} - \frac{39}{539}, \end{aligned}$$

and

$$\hat{X}_{11}^\perp = -\frac{94613488}{12589985808169}\delta^{2/3} - \frac{576368}{70074132513}\sqrt{33}\delta^{2/3} + \frac{140016}{290521} + \frac{12572512}{1912499743}\delta^{1/3}$$

$$-\frac{7072}{10644711}\sqrt{33}\delta^{1/3}.$$

Let

$$\begin{aligned}\varrho &= 2930798268 + 375926628\sqrt{69}, \\ \varsigma &= \frac{36264501 + 15017961\sqrt{69}}{2}, \\ \varpi &= \frac{67600081649 + 19193007225\sqrt{69}}{2}.\end{aligned}$$

Then (6.2) is true when

$$\begin{aligned}\widehat{b}_{23}^\perp &= \frac{273985}{406847} + \frac{320}{1220541}\varrho^{1/3} + \frac{2170961680}{3121974955532447}\varrho^{2/3} - \frac{6160}{69062263209}\sqrt{69}\varrho^{2/3}, \\ \widehat{a}_{23}^\perp &= -\frac{72}{406847}\varsigma^{1/3} - \frac{1377522036}{9422440425203183}\varsigma^{2/3} + \frac{30564}{532672306247}\sqrt{69}\varsigma^{2/3} + \frac{32952}{406847}, \\ \widehat{X}_{23}^\perp &= -\frac{2265083835349913600}{149274128311095576116540667}\varpi^{2/3} - \frac{585358470400}{366904827394808309061}\sqrt{69}\varpi^{2/3} \\ &\quad + \frac{343131204352}{496573444227} + \frac{779266147299200}{8609620666987823847}\varpi^{1/3} - \frac{243548800}{21161814311001}\sqrt{69}\varpi^{1/3}.\end{aligned}$$

We end this section with the quartic analogue of Ramanujan's table for $f(\ell)$. Once again, we wish to show that we can compute $Q_\ell(q)$ even without working with Hauptmoduls.

In the following table, we will state the value of N and in each entry, set

$$\beta^\perp = \alpha^\perp(q^N).$$

Table 3: Table of identities for $Q_N(q)$

$N = 3$	
Let	
	$X = \frac{(\alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{8},$
and	
	$Y = \frac{1 - (\alpha^\perp \beta^\perp)^{1/4} - ((1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{8},$
then	
	$Q_3(q) = (-384XY + 48X)^2.$
$N = 5$	

Let

$$X = \frac{(\alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/2}}{64},$$

and

$$Y = \frac{1 - (\alpha^\perp \beta^\perp)^{1/2} - ((1 - \alpha^\perp)(1 - \beta^\perp))^{1/2}}{32},$$

then

$$Q_5(q) = 48^2 (400XY - 20X + Y^2).$$

$$N = 7$$

Let

$$X = \frac{(2^4 \alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/8}}{4},$$

and

$$Y = \frac{2 - (2^4 \alpha^\perp \beta^\perp)^{1/4} - (2^4 (1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{32},$$

then

$$Q_7(q) = \left(-48 \frac{X(12712X^3 - 672XY - 9 - 67X + 144Y - 16128X^2Y)}{256X - 144Y + 9} \right)^2.$$

$$N = 11$$

Let

$$X = \frac{(2^4 \alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{16},$$

and

$$Y = \frac{1 - (\alpha^\perp \beta^\perp)^{1/4} - ((1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{8},$$

then

$$\sqrt{Q_{11}(q)} = \frac{F(X, Y)}{G(X, Y)},$$

where

$$F(X, Y) = -24 (-1982XY - 21301X^2 + 197120X^2Y^2 - 247X + 49Y \\ + 7928XY^2 + 482944X^3 - 10448Y^4 - 49280X^2Y + 5224Y^3 - 849Y^2),$$

and

$$G(X, Y) = 408Y - 192Y^2 + 127744XY - 15968X - 49 + 512Y^3.$$

$$N = 23$$

Let

$$X = \frac{(2^4 \alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/8}}{4},$$

and

$$Y = \frac{1 - (\alpha^\perp \beta^\perp)^{1/4} - ((1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{16},$$

then

$$A(X, Y)Q_{23}(q) + B(X, Y)\sqrt{Q_{23}(q)} + C(X, Y) = 0,$$

where

$$A(X, Y) = -19712Y - 13629X + 1232,$$

$$\begin{aligned} B(X, Y) = & 33663168X^2Y^2 - 3028410X^4 + 434652X^2 + 146893824XY^3 - 53803X \\ & - 23802240Y^3 - 5780544X^3Y - 4207896X^2Y + 1219 - 85080Y \\ & + 95208960Y^4 + 2008456XY - 27542592XY^2 + 361284X^3 + 2168280Y^2, \end{aligned}$$

$$\begin{aligned} C(X, Y) = & 9911568XY - 61307616X^2Y - 36105216Y^3 + 303176064X^4Y \\ & - 197184X - 58512Y + 21574368X^3Y - 4114722816X^2Y^3 \\ & - 6879492096XY^4 + 1822584X^2 + 771510528X^2Y^2 - 18948504X^4 \\ & - 186784608XY^2 - 172594944X^3Y^2 + 1719873024XY^3 - 915701760Y^5 \\ & + 2298528Y^2 - 3754104X^3 + 286156800Y^4 + 73405512X^5. \end{aligned}$$

$$N = 19$$

Let

$$X = \frac{(2^4 \alpha^\perp \beta^\perp (1 - \alpha^\perp)(1 - \beta^\perp))^{1/4}}{16},$$

and

$$Y = \frac{1 - (\alpha^\perp \beta^\perp)^{1/2} - ((1 - \alpha^\perp)(1 - \beta^\perp))^{1/2}}{32},$$

then

$$Q_{19}(q) = \frac{F(X, Y)}{G(X, Y)},$$

where

$$\begin{aligned} F(X, Y) = & 40956997588992X^2Y^2 + 412498219256064X^3Y^2 - 2284740864XY^4 \\ & - 614115712979712X^2Y^3 - 2304Y^5 + 833540779008X^3 \\ & - 40792878406656X^3Y + 12629403648X^2 - 1187160754176X^2Y \\ & - 4363776XY^2 + 3377267380998144X^2Y^4 + 7121866781011968X^4Y^2 \end{aligned}$$

$$\begin{aligned}
& + 151520256XY^3 + 46080XY + 24892565151744X^4 \\
& + 367591651531776X^5 - 799667514206208X^4Y - 8983708211810304X^5Y \\
& + 373735303335936X^3Y^3 + 2456180700954624X^6, \\
G(X, Y) = & -9229XY^2 + 4977X^3 - Y^3 - 384X^2 + 832XY - 2403X^2Y - 18X.
\end{aligned}$$

Note that we include the example $N = 19$ to show that it is possible to derive a series for $1/\pi$ for $N = 19$ even if we do not have a Hauptmodul to work with. We emphasize here that the identity between $Q_{19}(q)$ and the corresponding functions X and Y is just one of the infinitely many such relations satisfied by these functions.

Acknowledgement. The second author would like to thank Professor C. Krattenthaler for his hospitality and for providing an excellent research environment during his stay at the Faculty of Mathematics, University of Vienna.

REFERENCES

- [1] B.C. Berndt, *Ramanujan's Notebooks Part III*, Springer-Verlag, New York, 1991.
- [2] B.C. Berndt, H.H. Chan and W.C. Liaw, On Ramanujan's quartic theory of elliptic functions. *J. Number Theory* **88** (2001), no. 1, 129–156.
- [3] B.C. Berndt, H.H. Chan and L.-C. Zhang, Ramanujan's singular moduli, *The Ramanujan J.* **1** (1997), 53–74.
- [4] J.M. Borwein and P.B. Borwein, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*, Wiley, New York, 1987.
- [5] J.M. Borwein and P.B. Borwein, A cubic counterpart of Jacobi's identity and the AGM. *Trans. Amer. Math. Soc.* **323** (1991), 691–701.
- [6] J.M. Borwein, P.B. Borwein and F.G. Garvan, Some cubic identities of Ramanujan. *Trans. Amer. Math. Soc.* **343** (1994), 35–47.
- [7] H.H. Chan, Ramanujan's class invariants and Watson's empirical process. *J. London Math. Soc.* **57** (1998), Ser. 2, 545–561.
- [8] H.H. Chan, S.H. Chan and Z.G. Liu, Domb's numbers and Ramanujan–Sato type series for $1/\pi$, *Adv. Math.* **186** (2004), 396–410.
- [9] H.H. Chan and T.G. Chua, An alternative transformation formula for the Dedekind η -function via the Chinese Remainder Theorem, *Int. J. Number Th.* **12** (2016), no. 2, 513–526.
- [10] H.H. Chan and W.C. Liaw, On Russell-type modular equations, *Canad. J. Math.* **52** (2000), no. 1, 31–46.
- [11] H.H. Chan, A.C.P. Gee and V. Tan, Cubic Singular Moduli, Ramanujan's class invariant λ_n and the explicit Shimura Reciprocity Law. *Pacific J. Math.* **208** (2003), no. 1, 23–37.
- [12] H.H. Chan and M.L. Lang, Ramanujan's modular equations and the Atkin-Lehner involutions, *Israel J. Math.* **103** (1998), 1–16.
- [13] H.H. Chan, W.C. Liaw and V. Tan, Ramanujan's class invariant λ_n and a new class of series for $1/\pi$. *J. London Math. Soc.* **64** (2001), no. 1, 93–106.
- [14] A.C.P. Gee, Class invariants by Shimura's reciprocity law, *J. Théor. Nombres Bordeaux* **11** (1999), 45–72.
- [15] A.C.P. Gee and P. Stevenhagen, Generating class fields using Shimura reciprocity, *Algorithmic number theory* (Portland OR, 1998), 441–453, *Lecture notes in Comput. Sci.* 1423, Springer, Berlin, 1998.
- [16] C.J.C. Jacobi, *Gesammelte Werke, Bd. 1*, Reimer, Berlin, 1891.
- [17] H. Rademacher, *Topics in Analytic Number Theory*, Springer, Berlin, 1973.

- [18] S. Ramanujan, Modular equations and approximations to π , Quart. J. Math. (Oxford) **45** (1914), 350–372.
- [19] R. Russell, On $\kappa\lambda - \kappa'\lambda'$ modular equations, Proc. London Math. Soc. **19** (1887), 90–111.
- [20] P. Stevenhagen, Hilbert’s 12th problem, complex multiplication and Shimura reciprocity, Class field theory-its centenary and prospect (Tokyo, 1998), 161–176, Adv. Stud. Pure Math. **30**, Math. Soc. Japan, Tokyo, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 358 LITTLE HALL, GAINESVILLE, FL 32611, USA

E-mail address: alexb@ufl.edu

URL: <http://people.clas.ufl.edu/alexb>

MAIN ADDRESS: DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, BLOCK S17, 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076, SINGAPORE

TEMPORARY ADDRESS: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA

E-mail address: matchh@nus.edu.sg

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA

E-mail address: michael.schlosser@univie.ac.at

URL: <http://www.mat.univie.ac.at/~schlosse>